

A CLASS OF 2-LOCAL FINITE SPECTRA WHICH ADMIT A v_2^1 -SELF-MAP

PRASIT BHATTACHARYA^{1,*} AND PHILIP EGGER²

CONTENTS

1. Introduction	1
Organization of the paper	5
Acknowledgments	5
2. A -module structures on $B(2)$	6
3. Toda's realization theorems	9
4. Realization of B_2	15
5. Nonuniqueness of Z	19
6. Existence of a v_2^1 -self-map of Z	24
Appendix A. An explicit A -module structure on Z	27
A.1. A -module definition file for Bruner's program	28
A.2. Ext charts produced by Bruner's program	31
References	32

ABSTRACT. At the prime 2, Behrens, Hill, Hopkins and Mahowald showed $M_2(1, 4)$ admits 32-periodic v_2 -self-map and more recently authors joint with Mahowald showed A_1 also admits 32-periodic v_2 -self-map. This leads to the question, whether there exists a finite 2-local complex with periodicity less than 32. This paper will answer the question by producing a class of finite 2-local spectra $\tilde{Z}$ all of which admit 1-periodic v_2 -self-map.

Keywords: stable homotopy, v_2 -periodicity

1. INTRODUCTION

Let $\mathcal{C}_0$ be the category of p -local finite spectra, where p is a fixed prime. A v_n -self-map of an object X of $\mathcal{C}_0$ is a self-map $v : \Sigma^t X \rightarrow X$ such that

$$K(n)_* v : K(n)_* X \longrightarrow K(n)_* X$$

is an isomorphism. Here $K(n)$ is the n -th Morava K -theory and it is well-known that $K(n)_*$ is the graded ring $\mathbb{F}_p[v_n, v_n^{-1}]$ with $|v_n| = 2p^n - 2$. Since $K(n)_*$ is a graded field (i.e. every nonzero homogeneous element has a multiplicative inverse),

¹DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, 106 HAYES-HEALY HALL, NOTRE DAME, IN 46556

¹TEL: +1(574) 631-7776

*CORRESPONDING AUTHOR

²DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, 235 MCALLISTER BUILDING, UNIVERSITY PARK, PA 16802

²TEL: +1(814)865-1123

E-mail addresses: ¹pbbhattac@nd.edu, ²pee3@psu.edu.

$K(n)_*X$ is a graded vector space over $K(n)_*$ and the isomorphism $K(n)_*v$, up to a change of basis, is multiplication by a nonzero element of $K(n)_*$. We say that a v_n -self-map v has periodicity k if k is the smallest integer such that $K(n)_*v$ induces multiplication by v_n^k (in which case $t = (2p^n - 2)k$). We will abusively refer to a v_n -self-map of periodicity k as a v_n^k -self-map.

In 1998, Hopkins and Smith showed in [HS98] that for every $n \geq 0$, $\mathcal{C}_n$, the category of $K(n-1)$ -acyclics, is a thick subcategory of $\mathcal{C}_0$ and the $\mathcal{C}_n$ form a sequence of thick subcategories

$$\mathcal{C}_0 \supset \mathcal{C}_1 \supset \mathcal{C}_2 \supset \cdots \supset \mathcal{C}_\infty$$

where $\mathcal{C}_\infty$ is the category of contractible spectra. We say that a p -local finite spectrum X is of type n if $X \in \mathcal{C}_n \setminus \mathcal{C}_{n+1}$. They also showed that

Theorem 1.1 (Hopkins-Smith). *Every p -local finite spectrum X of type n admits a v_n -self-map*

$$v : \Sigma^{k(2p^n-2)}X \longrightarrow X.$$

Moreover, the cofiber Cv is a spectrum of type $n+1$.

Not only does Theorem 1.1 show the existence of v_n -self-maps, but it also provides a recipe for constructing type n spectra. However, [HS98] does not shed any light on the minimal periodicity of such a v_n -self-map, except to establish that the minimal periodicity is always a power of p .

One of the key properties of a v_n -self-map $v : \Sigma^t X \rightarrow X$ is that the iterated compositions

$$v^{\circ r} := \Sigma^{(r-1)t}v \circ \cdots \circ \Sigma^t v \circ v : \Sigma^{rt}X \longrightarrow X$$

are homotopically nontrivial and potentially give us an infinite family of elements in the stable homotopy groups of spheres. Typically, $v^{\circ r}$ composed with the inclusion of bottom cell

$$\tau_r : S^{rt} \xrightarrow{\text{incl}} \Sigma^{rt}X \xrightarrow{v^{\circ r}} X$$

is nontrivial. Therefore, the map τ_r factorizes through some skeleton, say $X^{\langle n_r \rangle}$, such that the composite with the pinch map to a top-cell of $X^{\langle n_r \rangle}$

$$\sigma_r : S^{m_r} \xrightarrow{\tau_r} X^{\langle n_r \rangle} \xrightarrow{\text{pinch}} S^{n_r}$$

is a nontrivial element of $\pi_{rt-n_r}(S^0)$. The collection of such $\{\sigma_r : r > 0\}$ forms an infinite family. The smaller the periodicity of v , the smaller the gap in degree between successive elements in the family. Hence the interest is in

- finding the minimal periodicity of the v_n -self-map on a given type n finite spectrum, and
- finding finite p -local spectra whose v_n -self-maps have periodicity as low as possible.

Recall that $K(0) = H\mathbb{Q}$ and v_0 is just multiplication by p . The sphere spectrum S^0 is a type 0 spectrum which admits a v_0^1 -self-map. Since S^0 admits a v_0^1 -self-map, any type 0 spectrum admits a v_0^1 -self-map.

The search for v_n -self-maps gets increasingly complicated as n increases. First, we remind ourselves some of the standard notations used in the literature. The cofiber of the $v_0^{i_0}$ -self-map of S^0 , i.e. multiplication by p^{i_0} , is called the i_0 -th Moore

spectrum at the prime p and is denoted by $M_p(i_0)$. By Theorem 1.1, $M_p(i_0)$ must admit a v_1 -self-map of some periodicity and the cofiber of

$$v_1^{i_1} : \Sigma^{i_1(2p-2)} M_p(i_0) \longrightarrow M_p(i_0)$$

is denoted by $M_p(i_0, i_1)$. In general, the cofiber of

$$v_n^{i_n} : \Sigma^{i_n(2p^n-2)} M_p(i_0, \dots, i_{n-1}) \longrightarrow M_p(i_0, \dots, i_{n-1})$$

is denoted by $M_p(i_0, \dots, i_{n-1}, i_n)$ and called a generalized Moore spectrum. Often in the literature a generalized Moore spectrum $M_p(i_0, \dots, i_n)$ with $i_k = 1$ for $0 \leq k \leq n$ is called a Smith-Toda complex and is denoted by $V_p(n)$. Alternatively, one can define the spectrum $M_p(i_0, \dots, i_n)$ as a realization of the BP_* -comodule

$$BP_*/\langle v_0^{i_0}, v_1^{i_1}, \dots, v_n^{i_n} \rangle.$$

Generalized Moore spectra may not exist for all sequences $(i_0, \dots, i_n)$ and even if such a spectrum exists, it may not be unique. Toda [To71] showed that $V_p(1) = M_p(1, 1)$ exists for $p \geq 3$, $V_p(2) = M_p(1, 1, 1)$ exists for $p \geq 5$ and $V_p(3) = M_p(1, 1, 1, 1)$ exists for $p \geq 7$. In 1966, J.F. Adams proved in [A66] that $M_2(1)$ does not admit a v_1^1 -self-map, in fact the minimal periodicity of a v_1 -self-map on $M_2(1)$ is 4. Thus, $M_p(1, i)$ does not exist unless $i \equiv 0 \pmod{4}$. In 2003, Behrens and Pemmaraju [BP04] showed that $V_3(1) = M_3(1, 1)$ admits a v_2^9 -self-map. Therefore $M_3(1, 1, i)$ does not exist unless $i \equiv 0 \pmod{9}$. In 2008, Behrens, Hill, Hopkins and Mahowald [BHHM08] showed that the v_2 -self-map of $M_2(1, 4)$ has periodicity 32. Little is known about v_n -self-maps for $n \geq 3$ aside from the work of Toda mentioned above, and Nave's proof in [N10] of the nonexistence of $V_p(\frac{p+1}{2})$ for $p > 7$.

Instead of focusing on generalized Moore spectra one can also ask the following question:

Question 1. *For a fixed prime p , what is the type n spectrum whose v_n -self-map has the smallest periodicity?*

For instance, at the prime 2, we have seen that $M_2(1)$ does not admit a v_1^1 -self-map. However, it is known that $Y := M_2(1) \wedge C\eta$ admits eight v_1^1 -self-maps (see [DM81]). At the prime 3, Behrens and Pemmaraju [BP04] showed that $M_3(1, 1)$ admits a v_2^9 -self-map. However, they also proved that $M_3(1, 1) \wedge Y(2)$, where

$$Y(2) = S^0 \cup_{\alpha_1} S^4 \cup_{2\alpha_1} S^8,$$

admits a v_2^1 -self-map. Toda proved that $M_p(1, 1)$ admits a v_2 -self-map for $p \geq 5$ and that $M_p(1, 1, 1)$ admits a v_3^1 -self-map for $p \geq 7$.

Though $M_2(1, 4)$ had a v_2^{32} -self-map, the authors hoped that one of the cofibers of the v_1^1 -self-maps on Y might have a v_2 -self-map of periodicity less than 32. In joint work with Mark Mahowald (see [BEM]), we showed that all of them have v_2^{32} -self-maps. This led to the question of whether there exists any 2-local type 2 spectrum with a v_2 -self-map of periodicity less than 32. We answer this question in the affirmative by producing a class of finite spectra that admit v_2^1 -self-maps. The main purpose of this paper is to prove the following theorem.

Main Theorem 1. *There is a collection of 2-local type 2 spectra $\tilde{\mathcal{Z}}$, such that every $Z \in \tilde{\mathcal{Z}}$ admits a v_2 -self-map of periodicity 1.*

For the rest of the paper, we will work in stable homotopy category of 2-local spectra. Let A denote the mod 2 Steenrod algebra and $A(n)$ be the subalgebra of

A generated by $\{Sq^{2^i} : 1 \leq i \leq n\}$. Let Q_n for $n \geq 0$ be the n -th Milnor element in A which are iteratively constructed using the formula

$$Q_n = [Sq^{2^n}, Q_{n-1}] = Sq^{2^n} Q_{n-1} - Q_{n-1} Sq^{2^n}$$

where $Q_0 = Sq^1$. The Milnor element Q_n generates an exterior algebra $E(Q_n)$ as $Q_n^2 = 0$ and it commutes with any $a \in A(n)$. The exterior algebra $E(Q_n)$ is a normal subalgebra of $A(n)$ and the pushout

$$A(n) // E(Q_n) = A(n) \otimes_{E(Q_n)} \mathbb{F}_2 = \mathbb{F}_2 \otimes_{E(Q_n)} A(n)$$

is an $A(n)$ -module, in fact, it is an $A(n)$ -algebra. Let $B(n)$ denote the $A(n)$ -algebra $A(n) // E(Q_n)$ and

$$q_n : A(n) \longrightarrow B(n)$$

denote the “quotient” map.

Definition 1.2. The class $\tilde{\mathcal{Z}}$ is the collection of all spectra Z such that there is an isomorphism of $A(2)$ -modules

$$H^*(Z) \cong B(2).$$

Notation 1.3. Any A -module which restricts to $A(2)$ as an $A(2)$ -module will be denoted A_2 . Likewise, any A -module which restricts to $B(2)$ as an $A(2)$ -module will be denoted B_2 .

Definition 1.2 is motivated by the fact that the cohomology of the spectrum Y , which admits v_1^1 -self-map, is

$$H^*(Y) = B(1).$$

To show that the class $\tilde{\mathcal{Z}}$ is nonempty, we first enrich the $A(2)$ -module $B(2)$ to an A -module B_2 . We do this by enriching the $A(2)$ -module $A(2)$ to an A -module A_2 and taking B_2 to be the image of q_2 . As we will point out in Remark 2.10 and expand upon in the appendix, there are many different A -modules A_2 , thus there are potentially many different A -modules B_2 . We will then topologically realize the B_2 as cohomologies of spectra (see Theorem 3.2).

There is yet another way of obtaining spectra in the class $\tilde{\mathcal{Z}}$. It is well known [HM14, Lemma 6.1] that there exists a nontrivial self-map

$$\gamma : \Sigma^5 C\eta \wedge C\nu \longrightarrow C\eta \wedge C\nu,$$

where η and ν are the well-known Hopf maps in $\pi_*(S^0)$. The map γ has multiple lifts

$$w : \Sigma^5 A_1 \wedge C\nu \longrightarrow A_1 \wedge C\nu$$

whose cofibers Cw belong to the class $\tilde{\mathcal{Z}}$. This approach of producing $Z \in \tilde{\mathcal{Z}}$ is described in the second author’s doctoral thesis [E, Chapter 3]. The authors believe that any spectrum in the class $\tilde{\mathcal{Z}}$ can be obtained as a cofiber of such a degree 5 self-map of $A_1 \wedge C\nu$.

Given such a B_2 , we see that the spectra realizing it are not unique, even up to homotopy. Depending on the specific B_2 we choose, there are either 4 or 8 different homotopy classes of spectra that realize B_2 (see Theorem 5.5).

Let $k(n)$ denote the connected cover of the n -th Morava K -theory. Lellmann [L82] proved that

$$H^*(k(n)) \cong A // E(Q_n).$$

Hopkins and Mahowald [HM14] showed that as an A -module

$$H^*(tmf) \cong A//A(2).$$

Since every $Z \in \tilde{\mathcal{Z}}$ is a realization of $A(2)//E(Q_2)$, we have

$$H^*(tmf \wedge Z) \cong A//A(2) \otimes A(2)//E(Q_2) \cong A//E(Q_2) \cong H^*(k(2)).$$

As a result, any spectrum $Z \in \tilde{\mathcal{Z}}$ satisfies the relation

$$(1.4) \quad tmf \wedge Z \simeq k(2).$$

Thus Z can be thought of as the height 2 analogue of the spectrum Y because

$$ko \wedge Y \simeq k(1).$$

As discussed earlier, Y admits a v_1^1 -self-map. Main Theorem 1 produces a v_2^1 -self-map of Z which further extends the analogy between Y and Z .

Understanding the ko -resolution of Y (see [M81] and [M82]) results in the proof of the telescope conjecture at chromatic height 1 at the prime 2. By analogy, we hope that the tmf -resolution of Z will enable us to attack the telescope conjecture at chromatic height 2 at the prime 2. Indeed, the authors have completely computed the “easier” side of the telescope conjecture, namely $\pi_*(L_{K(2)}Z)$, for any $Z \in \tilde{\mathcal{Z}}$. Part of this computation also appears in [E], with a more detailed report to appear.

Organization of the paper. In Section 2 we show that every A -module B_2 is “half” of a corresponding A_2 , in that there is a short exact sequence of A -modules

$$0 \longrightarrow \Sigma^7 B_2 \longrightarrow A_2 \longrightarrow B_2 \longrightarrow 0.$$

In Section 3 we recall a criterion of Toda for realizing a given A -module as the cohomology of a spectrum, and give a proof of a more refined criterion. In the process we review the construction of Adams towers and dual Adams towers, which will be necessary in Section 5.

In Section 4 we show that all A -modules B_2 satisfy Toda’s criterion and can thus be topologically realized. However, in Section 5, we show that all of these realizations are non-unique; given an A -module B_2 , there are, up to homotopy, either 4 or 8 different spectra that realize it.

Finally, in Section 6 we complete the proof of Main Theorem 1.

We provide Appendix A to show how to obtain A -module structure on $B(2)$ in practice. We obtain an explicit A -module and display it in the format required by Bruner’s Ext program [Bru93]. We also display various Ext charts obtained by running this program.

ACKNOWLEDGMENTS

The authors would like to thank Mark Behrens, Paul Goerss and Mike Mandell for their invaluable assistance and encouragement throughout this project. We would like to thank Irina Bobkova for providing Example 2.2 to point out a flawed argument in our previous draft, and Nicolas Ricka for suggestion Condition 2.4. We are indebted to Bob Bruner for his Ext calculator program. While the results in this paper are independent of any computer assisted programs, many of the insights in this paper have come about by analysing the outputs of this program. We are also grateful to Alex Kruckman for putting up an online program Adem relations calculator, which was very handy for our purposes.

2. A -MODULE STRUCTURES ON $B(2)$

Let Q_n be the Milnor element of A , let M be a left $A(m)$ -module for $m \geq n$ and let

$$\begin{aligned} \mathcal{Q}_n^R : \Sigma^{2^{n+1}-1} M &\rightarrow M \\ x &\mapsto xQ_n \end{aligned}$$

be the multiplication by Q_n on the right. Adams and Margolis [AM71] used the property $Q_n^2 = 0$ to define the Adams-Margolis homology

$$H(M; Q_n) = \frac{\ker \mathcal{Q}_n^R}{\text{img } \mathcal{Q}_n^R}$$

When $M = A(n)$, the right action of Q_n is same as the left action of Q_n as Q_n lies in the center of $A(n)$. Hence we can consider the map $\mathcal{Q}_n$ of multiplication by Q_n on the left or the right. Note that $\text{coker } \mathcal{Q}_n$ is isomorphic to $B(n)$. It can be easily checked that $H(A(n); Q_n) = 0$, therefore $\ker \mathcal{Q}_n \cong \text{img } \mathcal{Q}_n$. Moreover, the induced map $\tilde{\mathcal{Q}}_n$

$$\begin{array}{ccc} A(n) & \twoheadrightarrow & \text{coker } \mathcal{Q}_n \\ & \searrow & \downarrow \mathcal{Q}_n \\ & & \Sigma^{-7} \text{img } \mathcal{Q}_n \hookrightarrow \Sigma^{-7} A(n) \end{array}$$

$\tilde{\mathcal{Q}}_n \downarrow$

is in fact an isomorphism of $A(n)$ -modules. As a result (also see [Mit85]) we have the short exact sequence of $A(n)$ -modules

$$0 \longrightarrow \Sigma^{2^{n+1}-1} B(n) \cong \text{img } \mathcal{Q}_n \longrightarrow A(n) \xrightarrow{q_n} B(n) \cong \text{coker } \mathcal{Q}_n \longrightarrow 0.$$

Now we restrict our attention to $n = 2$ and consider the short exact sequence of $A(2)$ -modules

$$0 \longrightarrow \Sigma^7 B(2) \longrightarrow A(2) \xrightarrow{q_2} B(2) \longrightarrow 0.$$

Since this short exact sequence is not split, it corresponds to a nontrivial element

$$\tilde{v}_2 \in \text{Ext}_{A(2)}^{1,7}(B(2), B(2)).$$

Let B_2 denote an arbitrary left A -module whose underlying $A(2)$ -module structure is $B(2)$. In Theorem 2.6 we argue that $\tilde{v}_2$ lifts to an element

$$\bar{v}_2 \in \text{Ext}_A^{1,7}(B_2, B_2).$$


Thus there exists a short exact sequence of left A -modules

$$(2.1) \quad 0 \longrightarrow \Sigma^7 B_2 \xrightarrow{i_2} A_2 \xrightarrow{q_2} B_2 \longrightarrow 0,$$

where the underlying $A(2)$ -module structure of the A -module A_2 , is free over one generator in degree 0. This shows that any A -module structure on $B(2)$ can be obtained as $\text{img } \mathcal{Q}_2^R$ (or $\text{coker } \mathcal{Q}_2^R$) for a given A -module A_2 . In her thesis, Marilyn Roth showed that there exist 1600 different A -module structures on $A(2)$. Thus to obtain an A -module structure on $B(2)$ in practice, we should consider a left A -module structure A_2 on $A(2)$, and consider the A -modules $\text{img } \mathcal{Q}_2^R$ or $\text{coker } \mathcal{Q}_2^R$. But not every A -module structure on $A(2)$ will lead to an A -module structure on $B(2)$ as the underlying $A(2)$ -module structure of $\text{img } \mathcal{Q}_2^R$ (or $\text{coker } \mathcal{Q}_2^R$) may not be isomorphic to $B(2)$. We do not know of an explicit A -module structure on $A(2)$ for

which the underlying $A(2)$ -module structure on $\text{img } \mathcal{Q}_2^R$ differs from $B(2)$, however the following example shows that one cannot easily exclude the existence thereof.

Example 2.2. Consider the following left A -module structure on $A(1)$ which has 8 generators as a graded $\mathbb{F}_2$ -vector space.



The subscript on a generator indicates the degree of that generator. The red square brackets represent a left action of Sq^4 , the blue curved lines represent a left action of Sq^2 , and the black straight lines represent a left action of Sq^1 . In the category of $A(1)$ -modules, $\mathcal{Q}_0^R$ sends

$$\begin{aligned} g_0 &\mapsto g_1 \\ g_2 &\mapsto h_3 \\ g_3 &\mapsto h_4 \\ h_5 &\mapsto h_6 \end{aligned}$$

So as an $A(1)$ -module, $\text{img } \mathcal{Q}_0^R$ consists of the generators g_1, h_3, h_4 and h_6 . However in the category of A -modules, we have $h_5 = Sq^4 Sq^1 \cdot 1$, $\mathcal{Q}_0^R(h_5) = 0$. Therefore $\text{img } \mathcal{Q}_0^R$ consists of g_1, h_3 and h_4 . Hence the A -module structure of $\text{img } \mathcal{Q}_0^R$ does not restrict to the $A(1)$ -module structure of $\text{img } \mathcal{Q}_0^R$.

In the following lemma we give a condition which guarantees that the underlying $A(2)$ -module structure on $\text{img } \mathcal{Q}_2^R$ and $\text{coker } \mathcal{Q}_2^R$ is precisely $B(2)$.

Lemma 2.3. *Let A_2 denote an A -module whose underlying $A(2)$ -module structure is simply $A(2)$. Then the underlying $A(2)$ -module structure of $\text{img } \mathcal{Q}_2^R$ is $B(2)$ if and only if A_2 satisfies*

$$(2.4) \quad Q_2 Sq^8 Q_2 \cdot 1 = 0,$$

where 1 is the generator in degree 0.

Proof. Let $\mathcal{Q}_2$ denote the multiplication by Q_2 map in the category of $A(2)$ -modules and let $\mathcal{Q}_2^R$ denote the right multiplication by Q_2 map in the category of A -modules. The underlying $A(2)$ -module structure on $\text{img } \mathcal{Q}_2^R$ and $\text{coker } \mathcal{Q}_2^R$ is precisely $B(2)$ unless there exists $n \geq 3$ such that $Sq^{2^n} Q_2 \cdot 1$ does not belong to $\text{img } \mathcal{Q}_2$. For dimensional reasons we only need to check the case when $n = 3$. Note that $Q_2 Sq^8 Q_2 \cdot 1 = 0$ if and only if

$$Sq^8 Q_2 \cdot 1 = Q_2 a \cdot 1 = a Q_2 \cdot 1$$

for some $a \in A(2)$ as $\ker \mathcal{Q}_2 \cong \text{img } \mathcal{Q}_2$ and the result follows. $\square$

Before proving Theorem 2.6, we indulge ourselves in some preliminary computations of certain *Ext* groups. Let $\mathcal{G}$ denote a basis for $B(2)$ as a graded $\mathbb{F}_2$ -vector space. Observe that, as an $E(Q_2)$ -module

$$B(2) \cong \bigoplus_{c \in \mathcal{G}} \Sigma^{|c|} \mathbb{F}_2,$$

therefore

$$DB(2) \otimes B(2) \cong \bigoplus_{c \in D\mathcal{G} \times \mathcal{G}} \Sigma^{|c|} \mathbb{F}_2,$$

where $D\mathcal{G}$ is the basis of $DB(2)$ dual to $\mathcal{G}$. Consequently,

$$Ext_{E(Q_2)}^{*,*}(B(2), B(2)) \cong \mathbb{F}_2[v_2] \otimes \left(\bigoplus_{c \in D\mathcal{G} \times \mathcal{G}} \Sigma^{|c|} \mathbb{F}_2 \right),$$

where v_2 is the image of the periodicity generator of $Ext_{E(Q_2)}^{*,*}(\mathbb{F}_2, \mathbb{F}_2)$ in bidegree $(s, t) = (1, 7)$ induced by the unit map

$$\iota : DB(2) \otimes B(2) \longrightarrow \mathbb{F}_2.$$

One can use a change of rings isomorphism to see that

$$Ext_{A(2)}^{*,*}(B(2), B(2)) \cong Ext_{E(Q_2)}^{*,*}(\mathbb{F}_2, B(2)) \cong \mathbb{F}_2[v_2] \otimes \left(\bigoplus_{c \in D\mathcal{G}} \Sigma^{|c|} \mathbb{F}_2 \right).$$

We summarize the above discussion with the following lemma.

Lemma 2.5. *Let $\mathcal{G}$ be a basis for $B(2)$ as a graded $\mathbb{F}_2$ -vector space, $D\mathcal{G}$ be the corresponding basis for the dual $DB(2)$ and $\iota_0 \in \mathcal{G}$ be the unique generator in degree 0. Then we have isomorphisms*

(i)

$$Ext_{E(Q_2)}^{*,*}(B(2), B(2)) \cong \mathbb{F}_2[v_2] \otimes \left(\bigoplus_{c \in D\mathcal{G} \times \mathcal{G}} \Sigma^{|c|} \mathbb{F}_2 \right),$$

(ii)

$$Ext_{A(2)}^{*,*}(B(2), B(2)) \cong \mathbb{F}_2[v_2] \otimes \left(\bigoplus_{c \in D\mathcal{G}} \Sigma^{|c|} \mathbb{F}_2 \right),$$

and the inclusion of $E(Q_2)$ into $A(2)$ induces the v_2 -linear map

$$\begin{aligned} l : Ext_{A(2)}^{s,t}(B(2), B(2)) &\rightarrow Ext_{E(Q_2)}^{s,t}(B(2), B(2)) \\ \bar{g} &\mapsto (\bar{g}, \iota_0), \end{aligned}$$

for every $\bar{g} \in D\mathcal{G}$.

Theorem 2.6. *Let B_2 denote any A -module which restricts as an $A(2)$ -module to $B(2)$. Then there exists an element $\bar{v}_2 \in Ext_A^{1,7}(B_2, B_2)$ which maps to $\tilde{v}_2$ under the map*

$$k : Ext_A^{s,t}(B_2, B_2) \longrightarrow Ext_{A(2)}^{s,t}(B(2), B(2)).$$

Proof. Consider a minimal free A -module resolution of $DB_2 \otimes B_2$,

$$(2.7) \quad \dots \longrightarrow F_i \xrightarrow{f_i} \dots \longrightarrow F_2 \xrightarrow{f_2} F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} DB_2 \otimes B_2.$$

A minimal resolution has the property that all the differentials in the sequence

$$Hom_A(F_0, \mathbb{F}_2) \longrightarrow Hom_A(F_1, \mathbb{F}_2) \longrightarrow Hom_A(F_2, \mathbb{F}_2) \longrightarrow \dots$$

are trivial. Thus,

$$(2.8) \quad Ext_A^{i,*}(B_2, B_2) \cong Hom_A^*(F_i, \mathbb{F}_2) \cong \mathbb{F}_2 \langle A\text{-module basis of } F_i \rangle.$$

The identity map $1_{B_2} : B_2 \rightarrow B_2$ generates a nontrivial element

$$x_{0,0} \in Ext_A^{0,0}(DB_2 \otimes B_2, \mathbb{F}_2),$$

which corresponds to a basis element of F_0 by (2.8).

Now consider a minimal $A(2)$ -module resolution of $DB(2) \otimes B(2)$,

$$(2.9) \quad \dots \longrightarrow G_i \xrightarrow{g_i} \dots \longrightarrow G_2 \xrightarrow{g_2} G_1 \xrightarrow{g_1} G_0 \xrightarrow{g_0} DB(2) \otimes B(2).$$

There is a basis element $y_{0,0} \in G_0$ which corresponds to the

$$1_{B(2)} \in \text{Ext}_{A(2)}^{0,0}(B(2), B(2))$$

such that $k(x_{0,0}) = y_{0,0}$. From the $A(2)$ -module structure of $B(2)$, it is clear that

$$g_0(a \cdot y_{0,0}) \neq 0$$

for any $a \in A(2)$ with $0 \leq |a| \leq 6$ and

$$g_0(Q_2 \cdot y_{0,0}) = 0.$$

Therefore there is a generator $y_{1,7} \in G_1$ such that

$$g_1(y_{1,7}) = Q_2 \cdot y_{0,0}$$

which correspond to $\tilde{v}_2$. Since, any $a \in A$ with $0 \leq |a| \leq 7$ belongs to $A(2)$, same assertion holds for the element $x_{0,0}$, i.e. there is a basis element $x_{1,7} \in F_1$ such that

$$g_1(x_{1,7}) = Q_2 \cdot x_{0,0}.$$

The generator $x_{1,7}$ will correspond to an element

$$\bar{v}_2 \in \text{Ext}_A^{1,7}(DB_2 \otimes B_2, \mathbb{F}_2)$$

with the desired property. $\square$

Remark 2.10. (Number of A -module structures on B_2) Since any A -module structure on $B(2)$ can be produced as a quotient of an A -module structure of $A(2)$ which satisfies Condition 2.4, one can in principle count the number of A -module structures on $B(2)$ using the results of Roth [Ro]. This method of counting is extremely tedious as there are 1600 different A -module structures on $A(2)$. Moreover, the number of A -module structures does not reflect deeper concepts, nor is it directly related to the purpose of this paper. Nonetheless, in the appendix we discuss in detail how to use Roth's work [Ro] to produce A -module structures on $B(2)$ and demonstrate it via an example. The authors would be curious to know if there is a more elegant method of counting A -module structures on $B(2)$.

3. TODA'S REALIZATION THEOREMS

The purpose of this section is to review Toda's criteria for realizing an A -module as the cohomology of a spectrum. In the process we will review how to build Adams towers and dual Adams towers, which are essential in Section 5.

Let M be any graded bounded below A -module. Toda [To71, Lemma 3.1] gave a criterion for the existence of a spectrum X which realizes M , i.e.,

$$H^*(X) = M.$$

Theorem 3.1 (Toda). *Let M be a graded A -module which is bounded below. If $M^n \neq 0$ implies*

$$\text{Ext}_A^{s,n+s-2}(M, \mathbb{F}_2) = 0 \text{ for } s \geq 3,$$

then there exists a spectrum X such that $H^*(X) = M$.

There is yet another realization theorem due to Toda which has not been published before.

Theorem 3.2 (Toda). *Let M be a graded A -module which is bounded below. If*

$$\text{Ext}_A^{s,s-2}(M, M) = 0 \text{ for } s \geq 3,$$

then there exists a spectrum X such that $H^*(X) = M$.

A proof of Theorem 3.2 can be found in notes of Haynes Miller [Mil]. Since the result is unpublished we will give a proof along the lines prescribed in [Mil]. First note that Theorem 3.2 is stronger than Theorem 3.1 in the sense that the realization criterion in Theorem 3.1 implies the realization criterion in Theorem 3.2. To see this consider the algebraic Atiyah-Hirzebruch spectral sequence

$$E_1^{s,t,n} := DM^n \otimes Ext_A^{s,t}(M, \mathbb{F}_2) \Rightarrow Ext_A^{s,t+n}(M, M).$$

If for every $s \geq 3$ and every n such that $M^n \neq 0$, we have $Ext_A^{s,n+s-2}(M, \mathbb{F}_2) = 0$, then it follows that for all $s \geq 3$ and all $n \in \mathbb{Z}$, we have

$$E_1^{s,n+s-2,-n} = DM^{-n} \otimes Ext_A^{s,n+s-2}(M, \mathbb{F}_2) = 0.$$

Thus, the Atiyah-Hirzebruch spectral sequence forces

$$Ext_A^{s,s-2}(M, M) = 0$$

for every $s \geq 3$.

The broad idea is to consider a free A -module resolution of M

$$\dots \xrightarrow{d^i} F_i \longrightarrow \dots \xrightarrow{d^1} F_1 \xrightarrow{d^0} F_0 \twoheadrightarrow M$$

and build a corresponding tower of spectra

$$\longrightarrow X_i \longrightarrow \dots \longrightarrow X_1 \longrightarrow X_0$$

often called the *dual Adams tower*, such that

$$H^*(\lim_{\leftarrow} X_i) \cong M.$$

Recall that the *Adams tower* for a spectrum X consists of spectra $\{\tilde{X}_r : r \geq 0\}$ with maps

$$\begin{array}{ccccccc} \tilde{X}_0 = X & \xleftarrow{\tilde{i}_0} & \tilde{X}_1 & \xleftarrow{\tilde{i}_1} & \tilde{X}_2 & \xleftarrow{\tilde{i}_2} & \dots \longleftarrow \tilde{X}_r \xleftarrow{\tilde{i}_r} \dots \\ \tilde{k}_0 \downarrow & \nearrow \tilde{s}_0 & \tilde{k}_1 \downarrow & \nearrow \tilde{s}_1 & \tilde{k}_2 \downarrow & \nearrow \tilde{s}_2 & \dots \nearrow \tilde{k}_r \downarrow \nearrow \tilde{s}_r \dots \\ \tilde{K}_0 & & \tilde{K}_1 & & \tilde{K}_2 & & \dots \tilde{K}_r & \dots \end{array}$$

such that

- $\tilde{K}_r$ is a generalized Eilenberg McLane spectrum (GEM),
- the sequence

$$\tilde{X}_{r+1} \xrightarrow{\tilde{i}_r} \tilde{X}_r \xrightarrow{\tilde{k}_r} \tilde{K}_r$$

is a cofibration,

- $\tilde{s}_r : \tilde{K}_r \rightarrow \tilde{X}_{r+1}$ is the connecting map of degree -1 of the above cofiber sequence, and
- the composite $\tilde{d}_r = \Sigma \tilde{k}_{r+1} \circ \tilde{s}_r : \tilde{K}_r \rightarrow \Sigma \tilde{X}_{r+1} \rightarrow \Sigma \tilde{K}_{r+1}$ induces the map

$$d^r : F_{r+1} := H^*(\Sigma^{r+1} \tilde{K}_{r+1}) \longrightarrow F_r := H^*(\Sigma^r \tilde{K}_r).$$

Let X_r be the cofiber in the cofiber sequence

$$\tilde{X}_r \longrightarrow X \longrightarrow X_r.$$

The map $\tilde{i}_r : \tilde{X}_{r+1} \rightarrow \tilde{X}_r$ induces a map i_r

$$\begin{array}{ccccc} \tilde{X}_{r+1} & \longrightarrow & X & \longrightarrow & X_{r+1} \\ \tilde{i}_r \downarrow & & \parallel & & \downarrow i_r \\ \tilde{X}_r & \longrightarrow & X & \longrightarrow & X_r \\ \tilde{k}_r \downarrow & & \downarrow & & \downarrow k_r \\ \tilde{K}_r & \longrightarrow & * & \longrightarrow & K_r \end{array}$$

such that

$$X_{r+1} \xrightarrow{i_r} X_r \xrightarrow{k_r} K_r$$

forms a cofiber sequence, where $K_r \simeq \Sigma \tilde{K}_r$. The collection $\{X_r : r \geq 0\}$ is the dual Adams tower. Adams showed that if X is a bounded below spectrum, then we have

$$\lim_{\leftarrow} \tilde{X}_i \simeq *.$$

Therefore, we also have

$$\lim_{\leftarrow} X_i \simeq X_p,$$

where X_p is the p -completion of X . Just like the Adams tower, the dual Adams tower of a spectrum X fits into the diagram

$$(3.3) \quad \begin{array}{ccccccc} X_0 = * & \xleftarrow{i_0} & X_1 & \xleftarrow{i_1} & X_2 & \xleftarrow{i_2} & \cdots \xleftarrow{i_r} X_r \xleftarrow{i_r} \cdots \\ k_0 \downarrow & \nearrow s_0 & \downarrow k_1 & \nearrow s_1 & \downarrow k_2 & \nearrow s_2 & \cdots \downarrow k_r \nearrow s_r \cdots \\ K_0 & & K_1 & & K_2 & & \cdots K_r \cdots \end{array}$$

where $s_i : K_i \rightarrow \Sigma X_{i+1}$ are the connecting maps of the fiber sequences

$$X_{i+1} \longrightarrow X_i \longrightarrow K_i$$

and the composite

$$k_{i+1} \circ s_i : K_i \longrightarrow K_{i+1}$$

induces the map $d^r : F_{r+1} \rightarrow F_r$.

Proof of Theorem 3.2. Consider a free A -module resolution of M

$$\cdots \xrightarrow{d^r} F_r \xrightarrow{d^{r-1}} \cdots \xrightarrow{d^1} F_1 \xrightarrow{d^0} F_0 \longrightarrow M.$$

Let K_r be the GEM such that $H^*(K_r) \cong \Sigma^{1-r} F_r$. We intend to build a dual Adams tower as in (3.3) corresponding to this free resolution. Using the condition

$$Ext_A^{s,s-2}(M, M) = 0$$

for $s \geq 3$, we will show that M splits off $H^*(X_r)$ via the maps p_r and t_r as displayed in the diagram

$$(3.4) \quad \begin{array}{ccccccc} & & M & \xrightarrow{\cong} & M & \xrightarrow{\cong} & M \cdots \\ & \nearrow t_1 & & \nearrow t_2 & & \nearrow t_3 & \cdots \\ H^*(X_1) & \xrightarrow{i_1^*} & H^*(X_2) & \xrightarrow{i_2^*} & H^*(X_3) & \xrightarrow{i_3^*} & \cdots \\ & \searrow p_2 & & \searrow p_3 & & & \end{array}$$

Let $X = \varprojlim X_r$. The above splitting will ensure

$$\operatorname{colim}_{\rightarrow} H^*(X_r) \cong M.$$

Case 1: $r = 0, 1$ and 2. The first few cases are straightforward. We choose $X_0 = *$. Since

$$X_1 \longrightarrow X_0 \longrightarrow K_0$$

is a fiber sequence, it is immediate that $X_1 = \Sigma^{-1}K_0$. Choose $k_1 = d_0$ and let X_2 be the fiber in the sequence

$$X_2 \longrightarrow X_1 \xrightarrow{k_1} K_1.$$

Now we use the fact that $d_1 \circ \Sigma^{-1}d_0 : \Sigma^{-2}K_0 \rightarrow K_2$ is trivial to construct the map k_2 in the diagram

$$\begin{array}{ccccc} \Sigma^{-1}X_1 & \xrightarrow{\Sigma^{-1}k_1} & \Sigma^{-1}K_1 & \longrightarrow & X_2 \longrightarrow X_1 \\ & & \downarrow d_1 & \swarrow k_2 & \\ & & K_2 & & \end{array}$$

Let X_3 be the fiber of k_2 :

$$X_3 \longrightarrow X_2 \xrightarrow{k_2} K_2,$$

and let t_1 be the projection map

$$H^*(X_1) \cong H^*(\Sigma^{-1}K_0) \cong F_0 \longrightarrow M.$$

Producing the map $k_3 : X_3 \rightarrow K_3$ and the splitting $t_2 : H^*(X_2) \rightarrow M$ is the first nontrivial step of an inductive argument.

Case 2: $r = 3$. The fiber sequence $X_2 \rightarrow X_1 \rightarrow K_1$ produces a long exact sequence

$$\dots \longleftarrow \Sigma^{-1}F_0 \xleftarrow{d^0} \Sigma^{-1}F_1 \longleftarrow H^*(X_2) \longleftarrow F_0 \xleftarrow{d^0} F_1 \longleftarrow \dots$$

Since the cokernel of the map $d^0 : F_1 \rightarrow F_0$ is M we have the exact sequence of the top row in the diagram

$$(3.5) \quad \begin{array}{ccccccc} 0 & \longrightarrow & M & \xrightarrow{p_2} & H^*(X_2) & \longrightarrow & \Sigma^{-1} \ker d^0 \longrightarrow 0 \\ & & \uparrow \swarrow & & \uparrow \swarrow & & \downarrow \\ & & \Sigma^{-1}F_4 & \xrightarrow{d^3} & \Sigma^{-1}F_3 & \xrightarrow{d^2} & \Sigma^{-1}F_2 \xrightarrow{d^1} \Sigma^{-1}F_1 \end{array}$$

Since $d^1 \circ d^2 = 0$ and the right vertical arrow of the above diagram is a monomorphism, the map $k_2^* \circ d^2$ factors through a map $c_2 : \Sigma^{-1}F_3 \rightarrow M$. Notice that

$$p_2 \circ c_2 \circ d^3 = k_2^* \circ d^2 \circ d^3 = 0$$

and p_2 is injective, hence

$$c_2 \circ d^3 = 0.$$

Therefore the map c_2 represents a class

$$\bar{c}_2 \in \operatorname{Ext}_A^{3,1}(M, M).$$

Since $\operatorname{Ext}_A^{3,1}(M, M) = 0$, c_2 is a coboundary, i.e. c_2 factors through d^2

$$c_2 = b_2 \circ d^2$$

and

$$k_2^* \circ d^2 = p_2 \circ b_2 \circ d^2.$$

So if we replace k_2^* with $k_2^* - p_2 \circ b_2$, which is exactly of the type of alteration we are allowed to make, we see that

$$k_2^* \circ d^2 = 0.$$

Since the target of the map k_2 is a GEM, the algebraic alteration of k_2^* can be realized topologically. Therefore, we have a map k_3 in the diagram

$$\begin{array}{ccccc} \Sigma^{-1}X_2 & \xrightarrow{k_2} & \Sigma^{-1}K_2 & \longrightarrow & X_3 \longrightarrow X_2 \\ & & \downarrow d_2 & \swarrow k_3 & \\ & & K_3 & & \end{array}$$

Define X_4 to be the fiber of the map k_3 . Since, $k_2^* \circ d^2 = 0$, k_2^* factors through d^1 by a map which we denoted by a_3 in (3.5). Consequently, the exact sequence of the top row in the diagram of (3.5) splits and we have

$$t_2 : H^*(X_2) \longrightarrow M.$$

Case 3: $r > 3$. Now inductively assume that we have constructed

- $k_{r-1} : X_{r-1} \rightarrow K_{r-1}$,
- X_r as the fiber of the map k_{r-1} , and
- A diagram of maps

$$(3.6) \quad \begin{array}{ccccccc} 0 & \longrightarrow & M & \xleftarrow{p_{r-1}} & H^*(X_{r-1}) & \longrightarrow & \Sigma^{-r+2} \ker d^{r-3} \longrightarrow 0 \\ & & & \swarrow t_{p-1} & \uparrow k_{r-1}^* & \nwarrow a_{r-1} & \downarrow \\ & & & & \Sigma^{-r+2} F_{r-1} & \xrightarrow{d^{r-2}} & \Sigma^{-r+2} F_{r-2} \end{array}$$

whose top row is split exact.

The fiber sequence

$$X_r \longrightarrow X_{r-1} \longrightarrow K_{r-1}$$

produces the horizontal exact sequence in the diagram

(3.7)

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & M & & \Sigma M & & \\ & & \downarrow & & \downarrow & & \\ & & \downarrow t_{p-1} & & \downarrow t_{p-1} & & \\ \Sigma^{-r+2} F_{r-1} & \xrightarrow{k_{r-1}^*} & H^*(X_{r-1}) & \xrightarrow{i_{r-1}^*} & H^*(X_r) & \xrightarrow{\delta_{r-1}^*} & \Sigma^{-r+1} F_{r-1} \xrightarrow{k_{r-1}^*} H^{*+1}(X_{r-1}) \\ \downarrow d^{r-2} & \nearrow a_{r-1} & \downarrow & & \downarrow d^{r-2} & \nearrow a_{r-1} & \downarrow \\ \Sigma^{-r+2} F_{r-2} & \longleftarrow & \Sigma^{-r+2} \ker d^{r-3} & & \Sigma^{-r+1} F_{r-2} & \longleftarrow & \Sigma^{-r+1} \ker d^{r-3} \\ & & \downarrow & & & & \downarrow \\ & & 0 & & 0. & & \end{array}$$

The vertical split exact sequences are the part of the assumptions for the inductive step. By a diagram chase in the above diagram we find that

- the image of the map

$$k_{r-1}^* : \Sigma^{-r+2} F_{r-1} \longrightarrow H^*(X_{r-1})$$

is $\Sigma^{-r+2} \ker d^{r-3}$, hence

$$\ker i_{r-1}^* \cong \operatorname{coker} k_{r-1}^* \cong M,$$

and,

- the kernel of the map

$$k_{r-1}^* : \Sigma^{-r+1} F_{r-1} \longrightarrow H^{*+1}(X_{r-1})$$

is isomorphic to $\ker d^{r-2}$, hence

$$\operatorname{img} \delta_{r-1}^* \cong \Sigma^{-r+1} \ker d^{r-2}.$$

Consequently, the top row in the diagram (3.8)

$$\begin{array}{ccccccc} 0 & \longrightarrow & M & \xrightarrow{p_r} & H^*(X_r) & \longrightarrow & \Sigma^{-r+3} \ker d^{r-2} \longrightarrow 0 \\ & & \uparrow \swarrow & & \uparrow \swarrow & & \downarrow \\ & & | c_r & \searrow b_r & \uparrow k_r & \searrow a_r & \\ \Sigma^{-r+1} F_{r+2} & \xrightarrow{d^{r+1}} & \Sigma^{-r+1} F_{r+1} & \xrightarrow{d^r} & \Sigma^{-r+1} F_r & \xrightarrow{d^{r-1}} & \Sigma^{-r+1} F_{r-1} \end{array}$$

is an exact sequence. The diagram in (3.8) is just the generalization of the diagram in (3.5). Therefore, one can make exactly the same arguments as in the case of $r = 3$, to conclude k_r factors through a map c_r . The map c_r is a cocycle and it represents a class

$$\bar{c}_r \in \operatorname{Ext}_A^{r+1, r-1}(M, M).$$

Since $\operatorname{Ext}_A^{r+1, r-1}(M, M) = 0$, c_r is also a coboundary, hence factors through d^r via a map b_r . Replacing k_r^* with $k_r^* - p_r \circ b_r$, we see that

$$d_r \circ k_r = 0$$

in the diagram

$$\begin{array}{ccccc} \Sigma^{-1} X_r & \xrightarrow{k_r} & \Sigma^{-1} K_r & \longrightarrow & X_{r+1} \longrightarrow X_r \\ & & \downarrow d_r & \nearrow k_{r+1} & \\ & & K_{r+1} & & \end{array}$$

Hence we have a lift $k_{r+1} : X_{r+1} \rightarrow K_{r+1}$. Define X_{r+2} to be the fiber of k_{r+1} . Since $k_r^* \circ d^r = 0$, k_r^* factors through d^{r-1} via the map a_r . Therefore, the top row of the diagram in Equation 3.8 splits and we have a map

$$t_r : H^*(X_r) \longrightarrow M.$$

Convergence. Let $X = \varprojlim \tilde{X}_r$. We still need to show that

$$H^*(\varprojlim \tilde{X}_r) = \varinjlim H^*(\tilde{X}_r)$$

in order to conclude $H^*(X) = M$. This is true when M is bounded below. The argument is standard and is known as Adams' Convergence Theorem in the literature (see [A58, Theorem 2.1] and [A74, Part III, Theorem 15.1] for details).

□

Now we briefly discuss some basic properties of the Adams tower and the dual Adams tower of a spectrum. Let X be a k -connected spectrum, i.e. $\pi_i(X) = 0$ for $i \leq k$. The Adams tower $\{\tilde{X}_r : r \geq 0\}$ and the dual Adams tower $\{X_r : r \geq 0\}$ of X is said to be *minimal* if it corresponds to a minimal free A -module resolution of $H^*(X)$. Because minimal free resolutions of $H^*(X)$ are nonunique, it follows that minimal Adams towers and dual Adams towers of X are also nonunique. It follows from the construction, that for a minimal Adams tower of X , we will have

$$(3.9) \quad \text{Ext}_A^{s,t}(H^*(\tilde{X}_r), H^*(Y)) \cong \begin{cases} \text{Ext}_A^{s+r,t}(H^*(X), H^*(Y)) & \text{for } s \geq k+1 \\ 0 & \text{for } s \leq k \end{cases}$$

for any spectrum Y . The above isomorphism is realized by the map

$$\tilde{X}_r \longrightarrow X.$$

Similarly, a minimal dual Adams tower satisfies

$$(3.10) \quad \text{Ext}_A^{s,t}(H^*(X_r), H^*(Y)) \cong \begin{cases} \text{Ext}_A^{s,t}(H^*(X), H^*(Y)) & \text{for } s \leq r-1+k \\ 0 & \text{otherwise} \end{cases}$$

for an arbitrary spectrum Y .

Suppose we have a map of spectra $f : X \rightarrow Y$, where X and Y are both bounded below. Then f induces a map between their Adams towers

$$\begin{array}{ccccccccc} * & \longrightarrow & \dots & \longrightarrow & \tilde{X}_3 & \longrightarrow & \tilde{X}_2 & \longrightarrow & \tilde{X}_1 & \longrightarrow & X \\ \parallel & & & & \downarrow \tilde{f}_3 & & \downarrow \tilde{f}_2 & & \downarrow \tilde{f}_1 & & \downarrow f \\ * & \longrightarrow & \dots & \longrightarrow & \tilde{Y}_3 & \longrightarrow & \tilde{Y}_2 & \longrightarrow & \tilde{Y}_1 & \longrightarrow & Y \end{array}$$

and their dual Adams towers

$$\begin{array}{ccccccccc} X & \longrightarrow & \dots & \longrightarrow & X_3 & \longrightarrow & X_2 & \longrightarrow & X_1 & \longrightarrow & * \\ \downarrow f & & & & \downarrow f_3 & & \downarrow f_2 & & \downarrow f_1 & & \parallel \\ Y & \longrightarrow & \dots & \longrightarrow & Y_3 & \longrightarrow & Y_2 & \longrightarrow & Y_1 & \longrightarrow & *. \end{array}$$

However, the collection of maps $\{\tilde{f}_i : i \geq 0\}$ and $\{f_i : i \geq 0\}$ may not be unique, even when the Adams tower and its dual are minimal.

4. REALIZATION OF B_2

Let B_2 denote a fixed A -module structure on the $A(2)$ -module $B(2)$. The main purpose of this section is to use Toda's realization theorem, Theorem 3.1, to conclude:

Theorem 4.1. *There exists a spectrum $Z \in \tilde{\mathcal{Z}}$ such that*

$$H^*(Z) \cong B_2$$

as an A -module.

For this we need to compute $\text{Ext}_A^{*,*}(B_2, \mathbb{F}_2)$. For any A -module M there is a Bousfield-Kan spectral sequence

$$E_1^{s,t,n} = \bigoplus_n \text{Ext}_{A(2)}^{s-n,t}(\overline{A//A(2)})^{\otimes n} \otimes M, \mathbb{F}_2 \Rightarrow \text{Ext}_A^{s,t}(M, \mathbb{F}_2)$$

which is also otherwise known as the algebraic- tmf spectral sequence (see [BHHM08, BEM]). We will abbreviate the name to ‘alg- tmf SS’ for the rest of the paper.

Notation 4.2. To save space we will abbreviate $Ext_{A(2)}^{*,*}(N, \mathbb{F}_2)$ to $Ext_{A(2)}^{*,*}(N)$ and $Ext_{E(Q_2)}^{*,*}(K, \mathbb{F}_2)$ to $Ext_{E(Q_2)}^{*,*}(K)$, where N is an $A(2)$ -module and K is an $E(Q_2)$ -module.

Warning 4.3. The name ‘algebraic- tmf spectral sequence’ is due to the fact that $H^*(tmf) = A//A(2)$. However, there are similar spectral sequences involving $A//A(n)$ for $n \geq 3$, which are not realizable topologically. We point this out so that readers are aware of the fact that the results in this paper do not rely on the theory of tmf per se, unlike some other results on v_2 -self-maps of finite complexes (such as the results of [BEM, BHHM08, BP04]).

In [DM83] (also see [BHHM08, §5]), it has been proved that as an $A(2)$ -module

$$A//A(2) \cong \bigoplus_{j \geq 0} \Sigma^{8j} N_1(j)$$

where $N_1(j)$ is the j -th Brown-Gitler module [G83]. As a result the E_1 -page of the alg- tmf SS can be expressed as

$$E_1^{s,t,n} = \bigoplus_{n \geq 0} \left(\bigoplus_{j_1 \geq 0, \dots, j_n \geq 0} Ext_{A(2)}^{s-n, 8(j_1 + \dots + j_n)}(N_1(j_1) \otimes \dots \otimes N_1(j_n) \otimes M) \right).$$

We will refer to s as the Adams filtration, t as the internal degree and n as the tmf -filtration. Thus, the d_r -differentials have tridegree $(1, 0, r)$.

In Figure 1, we provide a visual aid to assist the understanding of the E_1 -page of the alg- tmf SS. We encode the tmf -filtration using colors and express the spectral sequence in $(x, y) = (t - s, s)$ coordinates. We use black for $n = 0$, blue for $n = 1$, red for $n = 2$ and green for $n = 3$. We draw the symbol $\nearrow$ with a sequence of numbers $j_1 \dots j_k$ at $(t - s, s) = (8(j_1 + \dots + j_k) - k, k)$ to indicate that we must place a shifted copy of $Ext_{A(2)}^{*,*}(N_1(j_k) \otimes \dots \otimes N_1(j_1) \otimes M)$ at that bidegree. By doing so, we assemble all the potential contributors to $Ext_A^{s,t}(M)$ in the $(t - s, s)$ coordinate system. With this arrangement, where we denote different alg- tmf filtrations using different colors, any differential in the alg- tmf SS looks like an Adams d_1 -differential.




FIGURE 1. A convenient pictorial description of the E_1 -page of the alg- tmf SS

Now we estimate $Ext_A^{s,t}(B)$ using the *alg-tmf* SS. Since B_2 as an $A(2)$ -module is isomorphic to $A(2)//E(Q_2)$, we can apply a change of rings formula to see

$$(4.4) \quad Ext_{A(2)}^{*,*}(N_1(j_k) \otimes \cdots \otimes N_1(j_1) \otimes B_2) \cong Ext_{E(Q_2)}^{*,*}(N_1(j_k) \otimes \cdots \otimes N_1(j_1)).$$

As an $E(Q_2)$ -module

$$N_1(1) \cong E(Q_2) \oplus \Sigma^4 \mathbb{F}_2 \oplus \Sigma^6 \mathbb{F}_2$$

and

$$N_1(2) \cong E(Q_2) \oplus \bigoplus_{2 \leq i \leq 4} \Sigma^{2i} E(Q_2) \oplus \bigoplus_{5 \leq i \leq 7} \Sigma^{2i} \mathbb{F}_2.$$

Computation of these Ext -groups over $E(Q_2)$ is very tractable. Firstly, $E(Q_2)$ and $\mathbb{F}_2$ are the only irreducible $E(Q_2)$ -modules, which means that any $E(Q_2)$ -module M can be expressed as direct sums of shifted copies of $E(Q_2)$ and $\mathbb{F}_2$. Moreover, the fact that

- $\mathbb{F}_2 \otimes \mathbb{F}_2 \cong \mathbb{F}_2$,
- $\mathbb{F}_2 \otimes E(Q_2) \cong E(Q_2) \otimes \mathbb{F}_2 \cong E(Q_2)$, and,
- $E(Q_2) \otimes E(Q_2) \cong E(Q_2) \oplus \Sigma^7 E(Q_2)$,

allows us to compute the irreducible decomposition of the products $M \otimes N$ of two $E(Q_2)$ -module. Once we know the irreducible decomposition of an $E(Q_2)$ -module M , we can compute $Ext_{E(Q_2)}^{*,*}(M, \mathbb{F}_2)$ using the facts

- $Ext_{E(Q_2)}^{*,*}(E(Q_2)) \cong \mathbb{F}_2$ and
- $Ext_{E(Q_2)}^{*,*}(\mathbb{F}_2) \cong \mathbb{F}_2[v_2]$, where v_2 has bidegree $(s, t) = (1, 7)$.




FIGURE 2. The E_1 -page of the *alg-tmf* SS for B_2

In Figure 2 we compute the first 25 stems of the E_1 -page of the *alg-tmf* SS. Note that the E_1 -page is a module over $Ext_{E(Q_2)}^{*,*}(\mathbb{F}_2) = \mathbb{F}_2[v_2]$, hence admits a v_2 -action. We use a $\circ$ to denote an element on which v_2 acts trivially, otherwise we use a $\bullet$. We use dotted lines to indicate the v_2 -action. We will indicate the *tmf* filtration using the color code established in Figure 1.

Proof of Theorem 4.1. B_2 is nonzero in dimensions 0 through 16. By Theorem 3.1, it is enough to prove

$$Ext_A^{s, s+n-2}(M) = 0$$

for $s \geq 3$ for $0 \leq n \leq 16$, equivalently,

$$Ext_A^{s,t}(M) = 0$$

for $s \geq 3$ and $-2 \leq t - s \leq 14$. From, Figure 2 it is clear that in the E_1 -page of $\text{alg-}tmf$ SS $E_1^{s,t,n} = 0$ for $s \geq 3$ and $-2 \leq t - s \leq 14$ for all $n \in \mathbb{N}$, and hence the result follows. $\square$

While we have proved what we set out to prove in this section, we would like to justify the May names and the differential in stem 22 of Figure 2, as it will be crucial for the proof of our main theorem in Section 6. To do this, we must delve deeper into the $\text{alg-}tmf$ SS, identifying the elements by the names they would inherit from the May spectral sequence.

When V is a subalgebra of A then there is a quasi-isomorphism of cobar complexes

$$C^{*,*}(\mathbb{F}_2, A, V) \cong C^{*,*}(\mathbb{F}_2, A//V, \mathbb{F}_2).$$

Therefore by filtering $C^{*,*}(\mathbb{F}_2, A//V, \mathbb{F}_2)$ using the May filtration one obtains a May spectral sequence computing $Ext_A^{*,*}(V)$. Unfortunately, B_2 is not a subalgebra of A . However, we will make use of the fact that

- $A(2)$ is a subalgebra of A , therefore has a May spectral sequence

$${}^{May}E_1^{*,*,*} = \mathbb{F}_2[h_{i,j} : i + j \geq 4] \Rightarrow Ext_A(A_2)$$

- and the cofiber of the map

$$\bar{v}_2 : \Sigma^6 B_2 \longrightarrow B_2$$

is A_2 in the derived category of A -modules,

to assign May names to certain elements in the E_1 page of $\text{alg-}tmf$ SS for B_2 .

Notation 4.5. For any A -module M we will denote the E_1 -page of the $\text{alg-}tmf$ SS by $E_1^{s,t,n}[M]$.

Note that

$$E_1^{s,*,n}[A_2] := Ext_{A(2)}^{s-n,*}(A(2) \otimes \overline{A//A(2)}^{\otimes n}) \cong Ext_{\mathbb{F}_2}^{s-n,*}(\overline{A//A(2)}^{\otimes n})$$

which is isomorphic to

$$\overline{A//A(2)}^{\otimes n}$$

when $n = s$ and 0 otherwise. Thus $E_1^{s,*,n}[A_2]$ is concentrated in filtrations $s = n$ and is isomorphic to the cobar complex for A_2

$$E_1^{n,t,n}[A_2] \cong C^{n,t}(\mathbb{F}_2, A//A(2), \mathbb{F}_2).$$

Therefore we can assign May names to elements of $E_1^{s,t,n}[A_2]$ by reading their cobar names. From (4.4) it is clear that every element x in the E_1 page of the $\text{alg-}tmf$ SS for B_2 satisfies either $v_2^i \cdot x \neq 0$ for all $i > 0$ or $v_2^1 \cdot x = 0$. Therefore the map

$$\bar{q}_2 : E_1^{s,t,n}[B_2] \longrightarrow E_1^{s,t,n}[A_2],$$

induced by $q_2 : A_2 \rightarrow B_2$, injects when restricted to the subcomplex $E_1^{s,*,s}[B_2]$. Therefore we can assign May names to the those elements in $E_1^{s,t,n}[B_2]$ for which $s = n$.

Let ζ_i denote the conjugate of ξ_i in the dual Steenrod algebra A_* . The A -module

$$\Sigma^8 N_1(1)_* \subset (A//A(2))_* = \mathbb{F}_2[\zeta_1^8, \zeta_2^4, \zeta_3^2, \zeta_4, \zeta_5, \dots]$$

consists of elements $\{\zeta_1^8, \zeta_2^4, \zeta_3^2, \zeta_4\}$ (see [DM83, BHHM08] for details). The red bullet

$$\bullet \in E_1^{2,2+22,2}[B_2]$$

is the contribution of a generator in $Ext_{E(Q_2)}^{0,24}(\Sigma^8 N_1(1) \otimes \Sigma^8 N_1(1))$ which corresponds to the cobar element $[\zeta_2^4 | \zeta_2^4]$. Therefore its May name is $h_{2,2}^2$. Similarly, the green circle

$$\circ \in E_1^{3,3+21,3}$$

is the contribution of a generator in $Ext_{E(Q_2)}^{3-3,24}(\Sigma^8 N_1(1) \otimes \Sigma^8 N_1(1) \otimes \Sigma^8 N_1(1))$ which corresponds to $[\zeta_1^8 | \zeta_1^8 | \zeta_1^8]$, hence has the May name h_3^3 . By work of Tangora [Tan66], which is also exposed in [K96], one has

$$(4.6) \quad d_2(h_{2,2}^2) = h_3^3$$

in the May spectral sequence, which explains the differential in Figure 2.

5. NONUNIQUENESS OF Z

In this section we prove that the spectra $Z \in \tilde{\mathcal{Z}}$ realizing a given A -module B_2 are never unique, even up to homotopy. In fact, a given A -module B_2 can be realized by either 4 or 8 homotopically different spectra in $\tilde{\mathcal{Z}}$. To motivate the proof, we first give a sufficient condition for a spectrum X under which the A -module structure of $H^*(X)$ determines X uniquely up to homotopy.

Proposition 5.1. *Let X be any finite spectrum, and let M denote the A -module $H^*(X)$. If*

$$Ext_A^{s,s-1}(M, M) = 0$$

for every $s \geq 2$, then any spectrum Y for which $H^*(Y) \cong M$ as an A -module, is weakly equivalent to X .

Proof. Consider the Adams spectral sequence

$$E_2^{s,t} = Ext_A^{s,t}(H^*(Y), H^*(X)) \Rightarrow [X, Y]_{t-s}.$$

Since $H^*(X) \cong H^*(Y) \cong M$, the E_2 page is isomorphic to $Ext_A^{s,s-1}(M, M)$. Let g denote a generator in bidegree $(0,0)$ which corresponds to the isomorphism $H^*(X) \cong H^*(Y)$ of A -modules. Since $Ext_A^{r,r-1}(M, M) = 0$ for $r \geq 2$, it follows that

$$d_r(g) = 0.$$

Thus g realizes a topological map

$$\bar{g}: X \longrightarrow Y$$

which induces an isomorphism in cohomology. Therefore, by Whitehead's theorem X is weakly equivalent to Y . $\square$

Therefore, to address the uniqueness question we must compute $Ext_A^{r,r-1}(B_2, B_2)$ for $r \geq 2$. We will use the $\text{alg-}tmf$ SS to do so. As an $A(2)$ -module $B_2 \cong A(2) // E(Q_2)$, therefore a change of ring formula implies

$$Ext_{A(2)}^{*,*}(N_1(j_k) \otimes \cdots \otimes N_1(j_1) \otimes DB_2 \otimes B_2) \cong Ext_{E(Q_2)}^{*,*}(N_1(j_k) \otimes \cdots \otimes N_1(j_1) \otimes DB_2).$$

Since

$$DB(2) \cong \bigoplus_{c \in D\mathcal{G}} \Sigma^{|c|} \mathbb{F}_2$$

as an $E(Q_2)$ -module, we have

$$\begin{aligned}
& E_1^{s,t,n}[DB_2 \otimes B_2] \\
& \downarrow \cong \\
& \bigoplus_{n \geq 0} \bigoplus_{j_1 \geq 0, \dots, j_n \geq 0} Ext_{E(Q_2)}^{s-n, 8(j_1 + \dots + j_n)}(N_1(j_1) \otimes \dots \otimes N_1(j_n) \otimes DB_2)) \\
& \downarrow \cong \\
& \bigoplus_{n \geq 0} \bigoplus_{j_1 \geq 0, \dots, j_n \geq 0} Ext_{E(Q_2)}^{s-n, 8(j_1 + \dots + j_n)}(N_1(j_1) \otimes \dots \otimes N_1(j_n) \otimes \bigoplus_{c \in DG} \Sigma^{|c|} \mathbb{F}_2)) \\
& \downarrow \cong \\
& \bigoplus_{c \in DG} \left(\bigoplus_{n \geq 0} \bigoplus_{j_1 \geq 0, \dots, j_n \geq 0} Ext_{E(Q_2)}^{s-n, 8(j_1 + \dots + j_n) + |c|}(N_1(j_1) \otimes \dots \otimes N_1(j_n)) \right) \\
& \downarrow \cong \\
& \bigoplus_{c \in DG} E_1^{s, t+|c|, n}[B_2].
\end{aligned}$$

In other words, $E_1^{*,*,*}[DB_2 \otimes B_2]$ is a direct sum of shifted copies of $E_1^{*,*,*}[B_2]$, one for each generator of DB_2 . We computed $E_1^{*,*,*}[B_2]$ in Section 4 and displayed it in Figure 2.




FIGURE 3. The E_1 -page of the algebraic- tmf spectral sequence for $DB_2 \otimes B_2$

Notation 5.2. We know that $H_*(Z) \cong \mathbb{F}_2[\xi_1, \xi_2] / \langle \xi_1^8, \xi_2^4 \rangle$ as an $A(2)_*$ -comodule. Let $g_{\xi_1^{i_1} \xi_2^{i_2}}$ be the element in $H^*(Z)$ dual to $\xi_1^{i_1} \xi_2^{i_2}$. We conveniently choose

$$\mathcal{G} = \{g_{\xi_1^{i_1} \xi_2^{i_2}} : 0 \leq i_1 \leq 7, 0 \leq i_2 \leq 3\}.$$

We denote the element of $D\mathcal{G}$, which is Spanier-Whitehead dual to $g_{\xi_1^{i_1} \xi_2^{i_2}}$, by $\bar{g}_{\xi_1^{i_1} \xi_2^{i_2}}$.

In Figure 3 we draw the alg- tmf SS for $DB_2 \otimes B_2$ for $0 \leq s \leq 3$ and $-2 \leq t - s \leq 2$. We place a $*$ in bidegree (s, t) if $E_1^{s,t,*}[DB_2 \otimes B_2] \neq 0$ but irrelevant to this discussion. In bidegree $(s, t) = (2, 2 - 1)$, the two bullets ($\bullet$) are $v_2^2 \cdot \bar{g}_{\xi_1^7 \xi_2^2}$ and $v_2^2 \cdot \bar{g}_{\xi_1^4 \xi_2^3}$, and the circle ($\circ$) which is $h_3^2 \cdot \bar{g}_{\xi_1^6 \xi_2^3}$.

Lemma 5.3. *Let B_2 denote any A -module structure on $B(2)$. Then*

$$\dim_{\mathbb{F}_2}(Ext_A^{2,1}(B_2, B_2)) = 2 \text{ or } 3$$

depending on the A -module structure on B_2 .

Proof. The elements $\{v_2^2 \cdot \bar{g}_{\xi_1^7 \xi_2^2}, v_2^2 \cdot \bar{g}_{\xi_1^4 \xi_2^3}, h_3^2 \cdot \bar{g}_{\xi_1^6 \xi_2^3}\}$ cannot support a nontrivial differential in alg- tmf SS as

$$E_1^{s,s-2,*}[DB_2 \otimes B_2] = 0$$

for all $s = 3$. Moreover, $v_2^2 \cdot \bar{g}_{\xi_1^7 \xi_2^2}$ and $v_2^2 \cdot \bar{g}_{\xi_1^4 \xi_2^3}$ cannot be a target of a differential in alg- tmf SS, as they are in algebraic- tmf filtration zero. Therefore, $v_2^2 \cdot \bar{g}_{\xi_1^7 \xi_2^2}$ and $v_2^2 \cdot \bar{g}_{\xi_1^4 \xi_2^3}$ are present in $Ext_A^{2,1}(B_2, B_2)$. However, $h_3^2 \cdot \bar{g}_{\xi_1^6 \xi_2^3}$ can be a target of a differential in alg- tmf SS for B_2 . $\square$

Remark 5.4. In fact, for the specific A -module structure on $B(2)$ which is worked out in Appendix A (see Figure 6), we see that

$$\dim_{\mathbb{F}_2}(Ext_A^{2,1}(B_2, B_2)) = 2$$

which means that $h_3^2 \cdot \bar{g}_{\xi_1^6 \xi_2^3}$ is trivial in the $Ext_A^{2,1}(B_2, B_2)$.

Theorem 5.5. *For an A -module B_2 whose underlying $A(2)$ -module is isomorphic to $B(2)$, let*

$$n = \dim_{\mathbb{F}_2}(Ext_A^{2,1}(B_2, B_2)).$$

Then there are 2^n different homotopy type of spectra $Z \in \tilde{\mathcal{Z}}$ such that

$$H^*(Z) \cong B_2$$

as an A -module.

Proof. By Theorem 3.2, we know that there exists at least one spectrum Z that realizes B_2 . Fix a minimal dual Adams tower $\{Z_i : i \geq 0\}$ for Z . The dual Adams tower for Z fits into the diagram

$$\begin{array}{ccccccc} Z_0 = * & \xleftarrow{i_0} & Z_1 & \xleftarrow{i_1} & Z_2 & \xleftarrow{i_2} & \dots \xleftarrow{i_r} Z_r \xleftarrow{i_r} \dots \\ \downarrow k_0 & \nearrow s_0 & \downarrow k_1 & \nearrow s_1 & \downarrow k_2 & \nearrow s_2 & \dots \downarrow k_r \nearrow s_r \dots \\ K_0 & & K_1 & & K_2 & & \dots & K_r & \dots \end{array}$$

where K_r is a GEM. The spectrum Z is then the limit

$$Z = \varprojlim Z_i.$$

To create another spectrum Y realizing B_2 , we alter k_2 using a nonzero element $\bar{\delta} \in Ext_A^{2,1}(B_2, B_2)$ in a way that the composite

$$d_r : K_r \xrightarrow{s_r} \Sigma Z_{r+1} \xrightarrow{k_{r+1}} \Sigma K_{r+1}$$

remains fixed for all $r \geq 0$. Then we argue that Y and Z are not weakly equivalent.

For an element $\bar{\delta} \in \text{Ext}_A^{2,1}(B_2, B_2)$ choose a cocycle representative

$$\delta^* : \Sigma F_2^* \longrightarrow B_2.$$

Since δ^* is a cocycle, the composite

$$\Sigma F_3^* \xrightarrow{d^2} \Sigma F_2^* \xrightarrow{\delta^*} M$$

is trivial. Note that $Z_1 = \Sigma^{-1}K_0$ and $k_1 = d_0$, therefore unwinding the long exact sequence associated to the fiber sequence

$$Z_2 \xrightarrow{i_1} Z_1 \xrightarrow{k_1} K_1$$

gives us the diagram

$$(5.6) \quad \begin{array}{ccccccc} \Sigma F_2^* & \xrightarrow{\Delta^*} & F_0^* \cong H^*(Z_1) & & & & \\ & \searrow \delta^* & \downarrow & \searrow i_1^* & & & \\ 0 & \longrightarrow & M & \xrightarrow{p_2} & H^*(Z_2) & \longrightarrow & \Sigma^{-1} \ker d^0 \longrightarrow 0 \\ & & & & \searrow k_1^* & & \downarrow \\ & & & & & & H^{*+1}(K_1), \end{array}$$

where the horizontal row is exact (compare with (3.5)). Let Δ^* denote a lift of δ^* , which exists as ΣF_2^* is a free A -module.

Now we build a dual Adams tower $\{Y_i : i \geq 0\}$ which fits into the diagram

$$\begin{array}{ccccccc} Y_0 = * & \xleftarrow{j_0} & Y_1 & \xleftarrow{j_1} & Y_2 & \xleftarrow{j_2} & \dots \xleftarrow{j_r} Y_r \xleftarrow{j_r} \dots \\ \downarrow l_0 & \nearrow t_0 & \downarrow l_1 & \nearrow t_1 & \downarrow l_2 & \nearrow t_2 & \dots \downarrow l_r \nearrow t_r \dots \\ K_0 & & K_1 & & K_2 & & \dots & & K_r & & \dots \end{array}$$

as follows. Define $Y_i := Z_i$ for $0 \leq i \leq 2$ and let $l_i = k_i$ for $0 \leq i \leq 1$. Let

$$l_2 : Y_2 \longrightarrow K_2$$

be the map that classifies

$$\tilde{k}_2^* + p_2 \circ \delta^* : \Sigma F_2^* \longrightarrow H^*(\tilde{Z}_2).$$

The condition that δ^* is a cocycle guarantees that $d_3 \circ \tilde{l}_2 \simeq 0$. By construction of l_2 , we have

$$(5.7) \quad \begin{array}{ccc} Y_2 = Z_2 & \xrightarrow{j_1=i_1} & Y_1 = Z_1 \\ \downarrow l_2 - k_2 & \nearrow \Delta & \\ & & K_2. \end{array}$$

Note that $\Delta \neq 0$ as $\bar{\delta} \neq 0$. However, $(l_2 - k_2) \circ t_1 = \Delta \circ j_1 \circ t_1 = 0$ which means

$$l_2 \circ t_1 = k_2 \circ s_1 = d_1.$$

Build the rest of the tower $\{Y_i\}$ in the usual way described in the proof of Theorem 3.2, and let Y denote the limit

$$Y := \varprojlim Y_i.$$

The choices for l_i for $i \geq 3$ will not make any difference as

$$Ext_A^{s,s-1}(B_2, B_2) = 0$$

for $s \geq 3$.

Note that

$$Hom_A(H^*(Y), H^*(Z)) \cong Ext_A^{0,0}(H^*(Y), H^*(Z)) = Ext_A^{0,0}(B_2, B_2) = \mathbb{F}_2.$$

Therefore there is exactly one A -module map $\iota_* : H^*(Y) \rightarrow H^*(Z)$ which is an isomorphism. Let $\bar{\iota}$ denote the corresponding element in $Ext_A^{0,0}(H^*(Y), H^*(Z))$. To show that Y and Z are not weakly equivalent it suffices to show that $\bar{\iota}$ is not a permanent cycle in Adams spectral sequence

$$(5.8) \quad E_2^{s,t} = Ext_A^{s,t}(H^*(Y), H^*(Z)) \Rightarrow [Z, Y]_{t-s}.$$

Indeed, assume $\bar{\iota}$ is a permanent cycle. Then there will be a map of spectra

$$\iota : Z \longrightarrow Y$$

which induces ι_* on cohomology and we will have a map of dual Adams towers

$$\begin{array}{ccccccc} Z & \longrightarrow & \dots & \longrightarrow & Z_3 & \longrightarrow & Z_2 \longrightarrow Z_1 \\ \downarrow \iota & & & & \downarrow \iota_3 & & \downarrow \iota_2 \quad \downarrow \iota_1 \\ Y & \longrightarrow & \dots & \longrightarrow & Y_3 & \longrightarrow & Y_2 \longrightarrow Y_1 \end{array}$$

Note that ι_1 and ι_2 exist trivially as $Z_1 = Y_1$ and $Z_2 = Y_2$. However, the diagram

$$\begin{array}{ccccc} Z_3 & \longrightarrow & Z_2 & \xrightarrow{k_2} & K_2 \\ \downarrow \iota_3 & & \parallel & \circlearrowleft & \parallel \\ Y_3 & \longrightarrow & Y_2 & \xrightarrow{l_2} & K_2 \end{array}$$

shows that the map $\iota_3 : Z_3 \rightarrow Y_3$ exists if and only if the right square commutes because horizontal rows are cofiber sequences. But (5.7) implies that the right square does not commute. Therefore ι_3 does not exist and we have a d_2 -differential

$$d_2(\bar{\iota}) = \bar{\delta}$$

in the Adams spectral sequence (5.8), hence a contradiction.

Thus we get exactly one homotopy type of spectra realizing B_2 for each element in $Ext_A^{2,1}(B_2, B_2)$ and the result follows. $\square$

Remark 5.9. Let Z_1 and Z_2 denote two spectra such that

$$H^*(Z_1) \cong H^*(Z_2) \cong B_2.$$

From the arguments in the proof of Theorem 5.5, we see that in the Adams spectral sequence

$$E_2^{s,t} = Ext_A^{s,t}(H^*(Z_2), H^*(Z_1)) \Rightarrow [Z_1, Z_2]_{t-s}$$

the generator $\bar{\iota}$ in bidegree $(0, 0)$ supports a nontrivial d_2 differential unless $Z_1 \simeq Z_2$. Therefore the d_2 -differential in some sense ‘measures the difference’ in homotopy type between the spectra Z_1 and Z_2 .

6. EXISTENCE OF A v_2^1 -SELF-MAP OF Z

As usual, let B_2 denote an A -module whose underlying $A(2)$ -module structure is $B(2)$. By Theorem 5.5 and Lemma 5.3, we know that there are either 4 or 8 different homotopy types of spectra which realize B_2 . Let Z denote a spectrum of a specific homotopy type which realizes B_2 . In Figure 4 we lay out the E_1 page of the alg- tmf SS for $DZ \wedge Z$.




FIGURE 4. The E_1 -page of the algebraic- tmf spectral sequence for $DB_2 \otimes B_2$

As in Figure 1 we use colors to distinguish tmf -filtration, with $*$ in bidegrees which are nonzero but irrelevant to the discussion. In Theorem 2.6, we have established that $v_2 \cdot \bar{t}$ is a nonzero permanent cycle in the alg- tmf SS denoted by $\bar{v}_2 \in Ext_A^{1,7}(H^*(Z), H^*(Z))$.

Proof of Main Theorem 1. From Figure 4 it is clear that

$$Ext_A^{s,s+5}(H^*(Z), H^*(Z)) = 0$$

for all $s \geq 4$. The task of proving the Main Theorem 1 then boils down to eliminating the possibility of $\bar{v}_2$ supporting a nontrivial d_2 -differential in the Adams spectral sequence

$$(6.1) \quad E_2^{s,t} = Ext_A^{s,t}(H^*(Z), H^*(Z)) \Rightarrow [Z, Z]_{t-s}.$$

Figure 4 makes it clear that in the E_1 page of alg- tmf SS for $DZ \wedge Z$

$$E_1^{3,8,*} = \{v_2^3 \cdot \bar{g}_{\xi_1^7 \xi_2^2}, v_2^3 \cdot \bar{g}_{\xi_1^4 \xi_2^3}, h_3^3 \cdot \bar{g}_{\xi_1^7 \xi_2^3}\}.$$

The element $h_3^3 \cdot \bar{g}_{\xi_1^7 \xi_2^4}$, which has tmf -filtration 3, is a target of a differential in the alg- tmf SS due to (4.6), hence trivial in the E_2 page of (6.1). However, the elements $v_2^3 \cdot \bar{g}_{\xi_1^7 \xi_2^2}$ and $v_2^3 \cdot \bar{g}_{\xi_1^4 \xi_2^3}$ are nonzero permanent cycles in the alg- tmf SS as they are in tmf -filtration 0, hence represent nonzero elements in the E_2 page of (6.1).

Since $H^*(k(2)) = A//E(Q_2)$, the E_2 page of the Adams spectral sequence computing $k(2)_*(DZ \wedge Z)$ is

$$Ext_{E(Q_2)}^{*,*,*}(H^*(Z), H^*(Z)) \cong Ext_{E(Q_2)}^{*,*,*}(B(2), B(2))$$

which we have already computed (see Lemma 2.5). Now consider the map of spectral sequences

$$(6.2) \quad \begin{array}{ccc} Ext_A^{s,t}(H^*(Z), H^*(Z)) & \Longrightarrow & \pi_{t-s}(DZ \wedge Z) \\ \downarrow k_* & & \downarrow k \\ Ext_{A(2)}^{s,t}(H^*(Z), H^*(Z)) & \Longrightarrow & tmf_{t-s}(DZ \wedge Z) \\ \downarrow l_* & & \downarrow l \\ Ext_{E(Q_2)}^{s,t}(H^*(Z), H^*(Z)) & \Longrightarrow & k(2)_{t-s}(DZ \wedge Z) \end{array}$$

induced by the maps $S^0 \xrightarrow{k} tmf \xrightarrow{l} k(2)$. The elements $v_2^i \cdot \bar{l}$, $v_2^i \cdot \bar{g}_{\xi_1^7 \xi_2^2}$ and $v_2^i \cdot \bar{g}_{\xi_1^4 \xi_2^3}$ for $i \geq 0$ have nonzero image under the map k_* as they are in tmf -filtration zero. Moreover, they have nonzero image under the composite $(l \circ k)_*$ by Lemma 2.5. From Remark 5.9, it is clear that

$$d_2(\bar{l}) = 0$$

in the upper-most spectral sequence in (6.2). Therefore,

$$d_2(k_*(l_*(\bar{l}))) = 0.$$

Since $k(2)$ admits a v_2 -self-map, the differentials in the bottom-most spectral sequence of (6.2) are v_2 -linear, which means

$$d_2(v_2 \cdot k_*(l_*(\bar{l}))) = 0.$$

Now, suppose that

$$d_2(\bar{v}_2) = c_1 v_2^3 \cdot \bar{g}_{\xi_1^7 \xi_2^2} + c_2 v_2^3 \cdot \bar{g}_{\xi_1^4 \xi_2^3}$$

with $(c_1, c_2) \neq (0, 0)$. This would imply

$$d_2(v_2 \cdot k_*(l_*(\bar{l}))) \neq 0$$

as $v_2^3 \cdot \bar{g}_{\xi_1^7 \xi_2^2}$ and $v_2^3 \cdot \bar{g}_{\xi_1^4 \xi_2^3}$ have nontrivial images under the map $k_* \circ l_*$, which is a contradiction. Thus we can conclude

$$d_2(\bar{v}_2) = 0$$

and Z admits a v_2^1 -self-map. $\square$

Remark 6.3 (v_2 -maps). Now let Z_1 and Z_2 be spectra of different homotopy type realizing B_2 . Then

$$(6.4) \quad d_2(\bar{l}) = c_1 v_2^2 \cdot \bar{g}_{\xi_1^7 \xi_2^2} + c_2 v_2^2 \cdot \bar{g}_{\xi_1^4 \xi_2^3} + c_3 h_3^2 \cdot \bar{g}_{\xi_1^6 \xi_2^3}$$

where $c_i \in \mathbb{F}_2$ and at least one of the $c_i \neq 0$. If either c_1 or c_2 is not zero then, essentially using the argument in the proof of Main Theorem 1, we see that

$$d_2(\bar{v}_2) = c_1 v_2^3 \cdot \bar{g}_{\xi_1^7 \xi_2^2} + c_2 v_2^3 \cdot \bar{g}_{\xi_1^4 \xi_2^3} \neq 0.$$

In other words, there is no map

$$v : \Sigma^6 Z_1 \longrightarrow Z_2$$

which induces multiplication by v_2^1 on Morava K -theory. Now suppose that there exists an A -module structure on B_2 for which $h_3^2 \cdot \bar{g}_{\xi_1^6 \xi_2^3}$ represents a nonzero element

in the E_2 page of the Adams spectral sequence of (6.1). Then we can choose Z_1 and Z_2 such that $c_1 = 0$, $c_2 = 0$ and $c_3 \neq 0$ in (6.4) and we will have a ‘ v_2^1 -map’

$$v : \Sigma^6 Z_1 \longrightarrow Z_2$$

between two different realizations of B_2 . It remains an open question whether such an A -module exists.

APPENDIX A. AN EXPLICIT A -MODULE STRUCTURE ON Z

The $A(2)$ -module $B(2)$ is 32-dimensional as an $\mathbb{F}_2$ -vector space spread across from degree 0 to degree 16. Endowing $B(2)$ with an A -module structure is therefore not easy in practice and requires a systematic approach. In Section 2, we established that any A -module structure on $B(2)$ extends to an A -module structure on $A(2)$ (see Theorem 2.6). Hence, we can obtain all possible A -module structures on $B(2)$ from A -module structures on $A(2)$. Marilyn Roth [Ro] showed that there are 1600 A -module structures on $A(2)$. However, she did not list explicit A -module structures or A_* -comodule structures. Rather, she listed 1600 different s -maps of the form

$$s : A(2)_* \longrightarrow (A//A(2))_*$$

satisfying certain criteria, each of which leads to a different A -module structure on $A(2)$. The purpose of this Appendix is to review Roth's work and demonstrate via an example different A -module structures on $B(2)$ in practice. We will express the A -module in the format required by Bruner's Ext software [Bru93] and display the output of the program.

We begin by describing the recipe for converting an s -map into an A -module structure on $A(2)$ as prescribed in [Ro]. It is useful to dualize things, considering not A -modules, but A_* -comodules. Recall that the dual Steenrod algebra A_* is the polynomial algebra

$$A_* = \mathbb{F}_2[\xi_i : i \geq 1]$$

where the generator ξ_i is in degree $2^i - 1$. As $A(2) \subset A$ is a sub-Hopf algebra generated by Sq^1 , Sq^2 and Sq^4 , the dual $A(2)_*$ is the quotient algebra

$$A(2)_* := \mathbb{F}_2[\xi_1, \xi_2, \xi_3] / \langle \xi_1^8, \xi_2^4, \xi_3^2 \rangle$$

of A_* , and

$$(A//A(2))_* = \mathbb{F}_2[\xi_1^8, \xi_2^4, \xi_3^2, \xi_4, \xi_5, \dots].$$

Notice that as an $\mathbb{F}_2$ -vector space, $A(2)_*$ has generators in degrees $0 \leq t \leq 23$. Between degrees 0 and 23, $(A//A(2))_*$ as an $\mathbb{F}_2$ -vector space, has exactly one nonzero generator in degrees 8, 12, 14, 15, 20, 22 and 23. In degrees 8, 12, 14, 15, 20, 22 and 23, $A(2)_*$ has 3, 4, 4, 3, 2, 1 and 1 generators respectively. As a result there are 2^{18} different s -maps. An s -map can be uniquely extended to an $\bar{s}$ -map

$$\bar{s} : A_* \longrightarrow (A//A(2))_*$$

as $A_* = A(2)_* \otimes (A//A(2))_*$. Corresponding to each s -map, there is a j -map

$$j : A(2)_* \longrightarrow A_*$$

which is defined as follows. Let $\pi : A_* \rightarrow A(2)_*$ be the canonical projection and $i : (A//A(2))_* \rightarrow A_*$ be the canonical inclusion. Define the j -map inductively using the formula

$$(A.1) \quad j(a) = a + s(a) + \sum_i j(\pi(a'_i)) \cdot \bar{s}(i(a''_i))$$

where a'_i and a''_i are part of the formula for the coproduct on a

$$\psi(a) = a \otimes 1 + 1 \otimes a + \sum_i a'_i \otimes a''_i.$$

Roth proved that [Ro, Chapter III]

$$A(2)_* \xrightarrow{j} A_* \xrightarrow{\psi} A_* \otimes A_* \xrightarrow{A_* \otimes \pi} A_* \otimes A(2)_*$$

defines an A_* -comodule structure on $A(2)_*$ if and only if the composite

$$(A.2) \quad A(2)_* \xrightarrow{j} A_* \xrightarrow{\psi} A_* \otimes A_* \xrightarrow{\bar{s} \otimes \bar{s}} (A//A(2))_* \otimes (A//A(2))_*$$

sends $\xi_1^7 \xi_2^3 \xi_3 \mapsto 0$. Roth listed all the s -maps which satisfy the criteria (A.2), and found that there are 1600 such s -maps.

One can obtain the A -module structure on $A(2)$ simply by dualizing the A_* -comodule structure on $A(2)_*$, however, in practice, this is quite tedious. Instead, one can obtain the A -module structure on $A(2)$ directly from the j -map using the right action of the total squaring operation

$$Sq = \sum_{i \geq 0} Sq^i : A_* \longrightarrow A_*.$$

Note that for a left A -module M , its dual $\widehat{M} = \text{Hom}_{\mathbb{F}_2}(M, \mathbb{F}_2)$ admits a right action of A . In particular, the right action of A on its dual A_* is determined by the right action of the total squaring operation Sq on A_* . It is well-known that the right action of Sq is a ring homomorphism determined by the formula

$$(\xi_i)Sq = \xi_i + \xi_{i-1}.$$

Given a j -map we obtain the A -module structure on $A(2)$ as follows. We declare

$$Sq^i(x) = x_1 + \cdots + x_n$$

where $x, x_1, \dots, x_n \in A(2)$, if $(j(x_{i*}))Sq^i$ contains $j(x_*)$ as a summand. After obtaining the A -module structure on $A(2)$ one must check if it satisfies Condition 2.4. If so, one can easily get an A -module structure on $B(2)$ by considering the inclusion map i_2 or the quotient map q_2 of (2.1).

A.1. A -module definition file for Bruner's program. We now consider the sample j -map that Roth computed [Ro, Pg 30, Chapter III] and check that the resultant A -module structure on $A(2)$ satisfies Condition 2.4. We will now express the resulting A -module structure on $B(2)$ as a definition file for Bruner's program [Bru93]. The A -module structure is encoded in a text file named **Z** in a way that we will now describe. The first line of the text file **Z** consists of a positive integer n , the dimension of $B(2)$ as an $\mathbb{F}_2$ -vector space, whose basis elements we will call $g_0, \dots, g_{n-1}$. The second line consists of an ordered list of integers $d_0, \dots, d_{n-1}$, which are the respective degrees of the g_i . Every subsequent line in the text file describes a nontrivial action of some Sq^k on some generator g_i . For instance, if we have

$$Sq^k(g_i) = g_{j1} + \cdots + g_{jl},$$

we would encode this fact by writing the line

$$\text{i k l j1} \dots \text{j1}$$

followed by a new line. Every action not encoded by such a line is assumed to be trivial. The text file **Z** is as follows.

32

0 1 2 3 3 4 4 5 5 6 6 6 7 7 7 8 8 9 9 9 10 10 10 11 11 12 12 13 13 14
15 16

0 1 1 1
0 2 1 2
0 3 1 3
0 4 1 5
0 5 1 7
0 6 1 9
0 7 1 12
0 10 1 20
0 12 1 25
0 13 1 27
0 14 1 29

1 2 2 3 4
1 3 1 6
1 4 2 7 8
1 5 1 10
1 6 2 12 13
1 7 1 15
1 8 1 17
1 9 1 20
1 12 1 27
1 14 1 30
1 15 1 31

2 1 1 3
2 2 1 6
2 4 3 9 10 11
2 5 2 12 14
2 6 2 15 16
2 7 1 18
2 8 1 21
2 9 1 23
2 10 1 25
2 11 1 27
2 12 1 29

3 2 1 8
3 3 1 10
3 4 2 12 14
3 6 3 17 18 19
3 7 2 20 22
3 8 2 23 24
3 9 1 26
3 10 2 27 28
3 11 1 29
3 12 1 30
3 13 1 31

4 1 1 6	10 10 1 31	18 4 1 27
4 2 1 8		18 6 1 30
4 3 1 10	11 1 1 14	18 7 1 31
4 4 1 13	11 2 1 16	
4 5 1 15	11 3 1 18	19 1 1 22
4 6 1 17	11 4 1 21	19 2 1 24
4 7 1 20	11 5 1 23	19 3 1 26
4 12 1 30	11 6 1 25	19 4 1 28
4 13 1 31	11 7 1 27	19 5 1 29
	11 10 1 31	19 6 1 30
5 1 1 7		19 7 1 31
5 2 2 9 10	12 2 1 17	
5 3 1 12	12 3 1 20	20 4 1 29
5 4 1 16	12 4 1 23	20 6 1 31
5 5 1 18	12 6 2 27 28	
5 6 2 20 22	12 7 1 29	21 1 1 23
5 12 1 31	12 8 1 30	21 2 2 25 26
	12 9 1 31	21 3 1 27
6 2 1 10		21 6 1 31
6 4 1 15	13 1 1 15	
6 6 1 20	13 2 1 17	22 2 1 26
6 12 1 31	13 3 1 20	22 4 1 29
	13 4 1 24	22 6 1 31
7 2 2 12 13	13 5 1 26	
7 3 1 15		23 2 2 27 28
7 4 2 17 18	14 2 2 18 19	23 3 1 29
7 5 1 20	14 3 1 22	23 4 1 30
7 6 1 24	14 4 2 23 24	23 5 1 31
7 7 1 26	14 5 1 26	
	14 6 2 27 28	24 1 1 26
8 1 1 10	14 7 1 29	24 4 1 30
8 4 2 17 19	14 8 1 30	24 5 1 31
8 5 2 20 22	14 9 1 31	
8 6 1 24		25 1 1 27
8 7 1 26	15 2 1 20	25 2 1 29
8 8 1 28	15 4 1 26	25 4 1 31
8 9 1 29		
8 10 1 30	16 1 1 18	26 4 1 31
8 11 1 31	16 2 1 22	
	16 4 2 25 26	27 2 1 30
9 1 1 12	16 5 1 27	27 3 1 31
9 2 1 15	16 6 1 29	
9 4 2 20 21		28 1 1 29
9 5 1 23	17 1 1 20	28 2 1 30
9 6 2 25 26	17 4 1 28	28 3 1 31
9 7 1 27	17 5 1 29	
9 8 1 29	17 6 1 30	29 2 1 31
	17 7 1 31	
10 4 2 20 22		30 1 1 31
10 6 1 26	18 2 1 24	
10 8 1 29	18 3 1 26	

A.2. Ext charts produced by Bruner's program. Using Bruner's program, we compute $Ext_A^{*,*}(B_2, \mathbb{F}_2)$ (see Figure 5) where B_2 is the A -module structure on $B(2)$ that we computed above. Bruner's program has features computes the A -module




FIGURE 5. The file Z_1.pdf displaying $Ext_A^{s,t}(B_2, \mathbb{F}_2)$ for $0 \leq t - s \leq 22$.

structure for Spanier-Whitehead duals and tensor product of two A -modules. Using these features we produce an A -module definition file for $B_2 \otimes DB_2$ which we named ZDZ. This allows us to compute $Ext_A^{*,*}(B_2, B_2)$ (see Figure 6) easily using Bruner's program.




FIGURE 6. The file ZDZ_1.pdf displaying $Ext_A^{s,t}(B_2, B_2)$ for $-7 \leq t - s \leq 15$.

REFERENCES

- [A66] Adams, J.F.: *On the groups $J(X)$. IV*, Topology 5 (1966), 21-71.
- [A58] Adams, J.F.: *On the structure and applications of the Steenrod algebra*. Comment. Math. Helv. 32 1958 180-214.
- [A74] Adams, J.F.: *Stable homotopy and generalised homology*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, Ill.-London, 1974. x+373 pp.
- [AM71] Adams, J.F. and Margolis, H.R.: *Modules over the Steenrod algebra*; Topology 10 (1971), 271-282.
- [BHHM08] Behrens, Mark, Hill, Michael A., Hopkins, Michael J. and Mahowald, Mark E.: *On the existence of a v_2^{32} self-map on $M(1, 4)$ at the prime 2*; Homology, Homotopy and Applications, Vol.10, No. 3 (2008), 45-84.
- [BEM] Bhattacharya, P., Egger, P. and Mahowald, Mark E.: *On the periodic v_2 -self-map of A_1* ; preprint: <https://arxiv.org/pdf/1406.3297.pdf>
- [Bru93] Bruner, Robert R.: *Ext in the nineties*; Algebraic topology (Oaxtepec, 1991), Contemp. Math. 146, Amer. Math. Soc., Providence, RI, 1993, pp. 71-90.
- [BP04] Behrens, M., Pemmaraju, S.: *On the existence of the self map v_2^9 on the Smith-Toda complex $V(1)$ at the prime 3*, Homotopy theory: relations with algebraic geometry, group cohomology, and algebraic K -theory, Contemp. Math. 346, Amer. Math. Soc., Providence, RI, 2004, pp. 9-49.
- [DM81] Davis, Donald M. and Mahowald, Mark E.: *v_1 and v_2 periodicity*; Amer. J. Math. 103, No. 4 (1981), 615-659
- [DM83] Davis, Donald M. and Mahowald, Mark E.: *Ext over the subalgebra A_2 of the Steenrod algebra for stunted projective spaces*, in Current trends in algebraic topology, Part 1 (London, Ont., 1981), CMS Conf. Proc., vol. 2, Amer. Math. Soc., Providence, RI, 1982, pp. 297-342.
- [E] Egger, Philip: *Some computations in $K(2)$ -local stable homotopy theory at the prime 2*; Ph.D. thesis.
- [G83] Goerss, Paul G.: *Some results on Brown-Gitler spectra*; Ph.D. thesis, 1983
- [GJM86] Goerss, Paul G., Jones, John D. S. and Mahowald, Mark E.: *Some generalized Brown-Gitler spectra*; Trans. Am. Math. Soc., Vol. 294, No. 1 (Mar., 1986), 113-132
- [HM14] Hopkins, Michael J. and Mahowald, Mark E.: *From elliptic curves to homotopy theory*. Topological modular forms, 294319, Math. Surveys Monogr., 201, Amer. Math. Soc., Providence, RI, 2014.
- [HS98] Hopkins, Michael J. and Smith, Jeffrey H.: *Nilpotence and Stable Homotopy Theory II*; Annals of Mathematics, 148 (1998), 1-49.
- [L82] Lellmann, Wolfgang: *Connected Morava K -theories*; Mathematische Zeitschrift, 179 (1982), 387-399.
- [May64] May, J. Peter: *The Cohomology of Restricted Lie Algebras*; Ph.D. thesis, 1964
- [M81] Mahowald, Mark E.: *bo-resolutions*; Pacific J. Math. 92 (1981), 365-383
- [M82] Mahowald, Mark E.: *The Image of J in the EHP sequence*; Annals of Mathematics 116 (1982), 65-112
- [Mil] Miller H.: *Toda's realization Theorem*; <http://www-math.mit.edu/~hrm/papers/toda-realization.pdf>.
- [Mit85] Mitchell, Stephen A.: *Finite complexes with $A(n)$ -free cohomology*. Topology 24 (1985), no. 2, 227-246.
- [K96] Kochman, Stanley O.: *Bordism, Stable Homotopy, and Adams Spectral Sequences*; Fields Institute Monographs, 7. American Mathematical Society, Providence, RI, 1996. xiv+272 pp. ISBN: 0-8218-0600-9.
- [Nak72] Nakamura, Osamu: *On the squaring operations in the May spectral sequence*; Mem. Fac. Sci. Kyushu Univ. Ser. A 26, No. 2 (1972), 293-308
- [N10] Nave, Lee S.: *The Smith-Toda complex $V((p+1)/2)$ does not exist*; Ann. of Math. (2) 171 (2010), no. 1, 491-509.
- [Rav88] Ravenel, Douglas C.: *Complex Cobordism and Stable Homotopy Groups of Spheres*; Bulletin (New Series) of the American Mathematical Society 18, 1988
- [Ro] Roth, Marylin J.: *The cyclic module structures of the hopf subalgebra A_2 over the Steenrod algebra and their geometric realizations*; Ph.D. thesis.
- [Tan66] Tangora, Martin C.: *On the cohomology of the Steenrod algebra*; Ph.D. thesis, 1966

- [To71] Toda, Hirosi.: *On spectra realizing exterior parts of the Steenrod algebra*. Topology 10 1971 53-65.